

	HOLDING (PUNCTURED VIAL)			SYRINGE STABILITY					DILUTION STABILITY						
	Storage Temperature	Light Protection	Result	Concentration	Diluent Pack	Storage Temperature	Light Protection	Result	Product Concentration	Vehicle (diluted with)	Diluent Pack	Storage Temperature	Light Protection	Result	
Carboplatin 10 mg/ml Study Date: 04/02/14 (holding + syringe + dilution)	20 - 25 °C	No	14 days	10 mg/ml	PP Plastic Syringe (BD)	20 - 25 °C	No	14 days	0.3 mg/ml	NaCl 0.9%	PE Bottle	20 - 25 °C	No	failed	
										Glucose 5%	PE Bottle	20 - 25 °C	No	42 days	
									3 mg/ml	NaCl 0.9%	PE Bottle	20 - 25 °C	No	failed	
										Glucose 5%	PE Bottle	20 - 25 °C	No	63 days	
Cisplatin 1 mg/ml Study Date: 04/02/14 (holding + syringe + dilution)	20 - 25 °C	No	14 days	1 mg/ml	PP Plastic Syringe (BD)	20 - 25 °C	No	30 days	0.01 mg/ml	NaCl 0.9%	PE Bottle	20 - 25 °C	No	35 days	
									0.4 mg/ml	NaCl 0.9%	PE Bottle	20 - 25 °C	No	35 days	
Cytarabine 100 mg/ml Study Date: 30/05/14 (syringe + dilution) Study Date: 16/12/14 (holding)	20 - 25 °C	No	14 days	100 mg/ml	PP Plastic Syringe (BD)	20 - 25 °C	No	60 days	0.19 mg/ml	Glucose 5%	PVC Latex free Bag	20 - 25 °C	No	14 days	
										NaCl 0.9%	PVC Latex free Bag	20 - 25 °C	No	84 days	
									7.6 mg/ml	NaCl 0.9%	PVC Latex free Bag	20 - 25 °C	No	84 days	
Docetaxel 20 mg/ml Study Date: 30/05/14 (dilution) Study Date: 10/09/12 (holding)	20 - 25 °C	No	21 days	-	-	-	-	-	0.3 mg/ml	NaCl 0.9%	Non PVC Bag	2 - 8 °C	Yes	98 days	
										Dextrose 5%	Non PVC Bag	2 - 8 °C	Yes	98 days	
										NaCl 0.9%	Non PVC Bag	20 - 25 °C	No	24 hours	
									0.9 mg/ml	Dextrose 5%	Non PVC Bag	20 - 25 °C	No	24 hours	
Doxorubicin 2 mg/ml Study Date: 04/02/14 (syringe + dilution)	-	-	-	2 mg/ml	PP Plastic Syringe (BD)	2 - 8 °C	Yes	96 days	0.05 mg/ml	NaCl 0.9%	PE Bottle	2 - 8 °C	Yes	84 days	
										Dextrose 5%	PE Bottle	2 - 8 °C	Yes	84 days	
									1.6 mg/ml	NaCl 0.9%	PE Bottle	2 - 8 °C	Yes	84 days	
										Dextrose 5%	PE Bottle	2 - 8 °C	Yes	84 days	
Epirubicin 2 mg/ml Study Date: 27/02/14 (syringe + dilution) Date: 16/12/14 (holding)	2 - 8 °C	Yes	14 days	2 mg/ml	PP Plastic Syringe (BD)	2 - 8 °C	Yes	96 days	0.05 mg/ml	NaCl 0.9%	PE Bottle	2 - 8 °C	Yes	84 days	
										Dextrose 5%	PE Bottle	2 - 8 °C	Yes	84 days	
									1.6 mg/ml	NaCl 0.9%	PE Bottle	2 - 8 °C	Yes	84 days	
										Dextrose 5%	PE Bottle	2 - 8 °C	Yes	84 days	
Etoposide 20 mg/ml Study Date: 22/12/14 (dilution) Study Date: 07/11/14 (syringe + holding)	20 - 25 °C	No	28 days	0.4 mg/ml in NaCl 0.9%	BD 50 ml Precise Syringe	20 - 25 °C	No	24 hours	0.2 mg/ml	NaCl 0.9%	PE Bottle	2 - 8 °C	Yes	28 days	
					BD 50 ml Precise Syringe	20 - 25 °C	No	24 hours			Dextrose 5%	PE Bottle	2 - 8 °C	Yes	28 days
					BD 50 ml Precise Syringe	20 - 25 °C	No	24 hours			NaCl 0.9%	PE Bottle	20 - 25 °C	No	28 days
					BD 50 ml Precise Syringe	20 - 25 °C	No	24 hours			Dextrose 5%	PE Bottle	20 - 25 °C	No	28 days
Fludarabine 25 mg/ml Study Date: 28/01/15 (syringe & dilution) Study Date: 11/07/14 (holding)	2 - 8 °C	Yes	28 days	0.25 mg/ml in NaCl 0.9%	BD 20 ml PP Syringe	2 - 8 °C	Yes	115 days	0.25 mg/ml	NaCl 0.9%	PE Bag	2 - 8 °C	Yes	115 days	
					BD 20 ml PP Syringe	20 - 25 °C	No	115 days			NaCl 0.9%	PE Bag	20 - 25 °C	No	115 days
					BD 20 ml PP Syringe	2 - 8 °C	Yes	115 days			NaCl 0.9%	PE Bag	20 - 25 °C	No	115 days
					BD 20 ml PP Syringe	20 - 25 °C	No	115 days			Dextrose 5%	PE Bag	20 - 25 °C	No	115 days
Fluorouracil 50 mg/ml Study Date: 27/02/14 (syringe + dilution) Study Date: 16/08/13 (PUMP) Study Date: 13/11/12 (holding)	20 - 25 °C	No	7 days	50 mg/ml	PP Plastic Syringe (BD)	20 - 25 °C	No	96 days	0.1 mg/ml	NaCl 0.9%	PUMP	2 - 8 °C	Yes	28 days	
										Glucose 5%	PUMP	20 - 25 °C	No	28 days	
										NaCl 0.9%	PUMP	2 - 8 °C	Yes	28 days	
										Glucose 5%	PUMP	20 - 25 °C	No	28 days	
									0.5 mg/ml	NaCl 0.9%	Non PVC Bag	20 - 25 °C	No	112 days	
										Dextrose 5%	Non PVC Bag	20 - 25 °C	No	100 days	
										NaCl 0.9%	Non PVC Bag	20 - 25 °C	No	112 days	
										Dextrose 5%	Non PVC Bag	20 - 25 °C	No	100 days	
									10 mg/ml	NaCl 0.9%	PUMP	2 - 8 °C	Yes	28 days	
										NaCl 0.9%	PUMP	20 - 25 °C	No	28 days	
										Glucose 5%	PUMP	2 - 8 °C	Yes	28 days	
										Glucose 5%	PUMP	20 - 25 °C	No	28 days	
25 mg/ml	NaCl 0.9%	PUMP	2 - 8 °C	Yes	28 days										
	NaCl 0.9%	PUMP	20 - 25 °C	No	28 days										
	Glucose 5%	PUMP	2 - 8 °C	Yes	28 days										
	Glucose 5%	PUMP	20 - 25 °C	No	28 days										
50 mg/ml	NaCl 0.9%	PUMP	2 - 8 °C	No	28 days										
	NaCl 0.9%	PUMP	20 - 25 °C	No	28 days										
Gemcitabine Powder Study Date: 21/10/10 (holding + dilution)	20 - 25 °C	No	28 days	-	-	-	-	-	0.1 mg/ml	NaCl 0.9%	Non PVC Bag	20 - 25 °C	No	35 days	
									10 mg/ml	NaCl 0.9%	Non PVC Bag	20 - 25 °C	No	35 days	
Gemcitabine 100 mg/ml Study Date: 30/05/14 (dilution) Study Date: 20/08/14 (holding)	20 - 25 °C	No	28 days	-	-	-	-	-	0.1 mg/ml	NaCl 0.9%	Non PVC Bag	20 - 25 °C	No	84 days	
									10 mg/ml	NaCl 0.9%	Non PVC Bag	20 - 25 °C	No	84 days	
Methotrexate 25 mg/ml Study Date: 23/08/13 (holding + dilution)	20 - 25 °C	No	56 days	-	-	-	-	-	0.1 mg/ml	NaCl 0.9%	PE Bag	2 - 8 °C	Yes	56 days	
										Glucose 5%	PE Bag	20 - 25 °C	No	2 days	
										NaCl 0.9%	PE Bag	2 - 8 °C	Yes	56 days	
										Glucose 5%	PE Bag	20 - 25 °C	No	2 days	
									20 mg/ml	NaCl 0.9%	PE Bag	2 - 8 °C	Yes	56 days	
										Glucose 5%	PE Bag	20 - 25 °C	No	56 days	
Ondansetron 2mg/ml Study Date: 12/03/12 (holding) Study date: SPC (dilution)	20 - 25 °C	No	4 days	-	-	-	-	-	0.16 mg/ml	Mannitol 10%	PE Bag	2 - 8 °C	Yes	24 hours	
											Ringer	PE Bag	20 - 25 °C	No	24 hours
	20 - 25 °C	Yes	7 days							Potassium Chloride 0.3% + Glucose 5%	PE Bag	2 - 8 °C	Yes	24 hours	
										Potassium Chloride 0.3% + Glucose 5%	PE Bag	20 - 25 °C	No	24 hours	
Oxaliplatin 5 mg/ml Study Date: 25/05/12 (holding) Study Date: 14/08/13 (dilution)	20 - 25 °C	No	14 days	-	-	-	-	-	0.1 mg/ml	Glucose 5%	Non PVC Bag	20 - 25 °C	No	84 days	
									1.2 mg/ml	Glucose 5%	Non PVC Bag	20 - 25 °C	No	84 days	
Paclitaxel 6 mg/ml Study Date: 04/02/14 (holding + dilution) Study Date: 10/07/2012 (dilution in fridge + Dextrose & NaCl + Dextrose in Ringer Solution)	20 - 25 °C	No	14 days	-	-	-	-	-	0.1 mg/ml	NaCl 0.9%	PE Bottle	20 - 25 °C	No	21 days	
										Glucose 5%	PE Bottle	20 - 25 °C	No	21 days	
									0.3 mg/ml	NaCl 0.9%	PE Bottle	2 - 8 °C	Yes	14 days	
										Dextrose 5% in Ringer Solution	PE Bottle	20 - 25 °C	No	17 days	
											PE Bottle	2 - 8 °C	Yes	14 days	
											PE Bottle	20 - 25 °C	No	14 days	
											PE Bottle	2 - 8 °C	Yes	14 days	
											PE Bottle	20 - 25 °C	No	14 days	
									1.2 mg/ml	Glucose 5%	PE Bottle	20 - 25 °C	No	14 days	
										NaCl 0.9%	PE Bottle	2 - 8 °C	Yes	7 days	
											PE Bottle	20 - 25 °C	No	14 days	
										Dextrose 5% in Ringer Solution	PE Bottle	2 - 8 °C	Yes	14 days	
	PE Bottle	20 - 25 °C	No	14 days											
	PE Bottle	2 - 8 °C	Yes	14 days											
Topotecan Powder Study Date: 16/04/11 (holding + dilution)	2 - 8 °C	Yes	28 days	-	-	-	-	-	0.01 mg/ml	NaCl 0.9%	PVC Bottle	2 - 8 °C	Yes	28 days	
											Dextrose 5%	PVC Bottle	20 - 25 °C	No	7 days
	20 - 25 °C	No	28 days							NaCl 0.9%	PVC Bottle	2 - 8 °C	Yes	28 days	
										Dextrose 5%	PVC Bottle	20 - 25 °C	No	21 days	
Topotecan 1 mg/ml Study Date: 30/05/14 (holding + dilution) Date: 10/07/12 (dilution: 0.025 mg)	20 - 25 °C	No	30 days	-	-	-	-	-	0.025 mg/ml	NaCl 0.9%	Non PVC Bag	2 - 8 °C	No	30 days	
											Non PVC Bag	20 - 25 °C	No	30 days	
											NaCl 0.9%	Non PVC Bag	2 - 8 °C	No	30 days
											Non PVC Bag	20 - 25 °C	No	30 days	
										0.1 mg/ml	NaCl 0.9%	Non PVC Bag	20 - 25 °C	No	84 days
											Glucose 5%	Non PVC Bag	20 - 25 °C	No	84 days
0.5 mg/ml	NaCl 0.9%	Non PVC Bag	20 - 25 °C	No	84 days										
	Glucose 5%	Non PVC Bag	20 - 25 °C	No	84 days										